DP CHEM SUMMER ASSIGNMENT

Incoming 11th graders Summer 2022

- 1. <u>Due first day of class (in order to do lab)</u>: **Watch the safety video (safety video by American Chemical society)** and answer the Safety quiz (front and back) https://www.youtube.com/watch?v=9077QEeM-68
- 2. Complete the Full Summer Review. Show all work.

You will have a pre-test on MYP content and content on this assignment (including knowing polyatomic ions)

3. Remember: This is a big packet to make sure that you are prepared for DP chem. Do not wait until the last minute to complete the packet. I will be available to answer any questions the first week of school but you should be able to do most of it on your own as all this content was covered in MYP.

Name	Formula	Name	Formula	
Acetate	C ₂ H ₃ O ₂ ⁻	Hypochlorite	CIO-	
Carbonate	CO ₃ ²⁻	Chlorite	CIO2-	
Hydrogen carbonate (or bicarbonate)	HCO ₃	Chlorate	CIO ₃	
Hydroxide	OH-	Perchlorate	CIO ₄	
Nitrite	$N0_2^-$	Permanganate	MnO_4^-	
Nitrate	NO ₃	Sulfite	SO32-	
Chromate	CrO ₄ ²⁻	Hydrogen sulfite (or bisulfite)	HSO ₃	
Dichromate	Cr ₂ O ₇ ²⁻	Sulfate	SO ₄ ²⁻	
Phosphate	PO ₄ ³⁻	Hydrogen sulfate (or bisulfate)	${\rm HSO_4}^-$	
Hydrogen phosphate	HPO ₄ ²⁻	Cyanide	CN-	
Dihydrogen phosphate	$H_2PO_4^-$	Peroxide	022-	
Ammonium	NH ₄ ⁺			

Thiocyanate SCN⁻

NO₃⁻nitrate SO₄²⁻sulfate PO₄³⁻phosphate NO₂⁻nitrite SO₃²⁻sulfite PO₃³⁻ phosphite ClO⁻hypochlorite BrO⁻hypobromite IO⁻hypoiodite ClO₂⁻chlorite

BrO₂ bromite IO₂ iodite

ClO₃ chlorate BrO₃ bromate IO₃ iodate

ClO₄ *perchlorate* BrO₄ *perbromate* IO₄ *periodate*

Significant Figures in Measurement and Calculations

A successful chemistry student habitually labels all numbers, because the unit is important. Also of great importance is the number itself. Any number used in a calculation should contain only figures that are

considered reliable; otherwise, time and effort are wasted. Figures that are considered reliable are called significant figures. Chemical calculations involve numbers representing actual measurements. In a measurement, significant figures in a number consist of:

Figures (digits) definitely known + One estimated figure (digit)

In class you will hear this expressed as "all of the digits known for certain plus one that is a guess."

Recording Measurements

When one reads an instrument (ruler, thermometer, graduate, buret, barometer, balance), he expresses the reading as one which is reasonably reliable. For example, in the accompanying illustration, note the

reading marked A. This reading is definitely beyond the 7 cm mark and also beyond the 0.8 cm mark. We read the 7.8 with certainty. We further estimate that the reading is five-tenths the distance from the 7.8 mark to the 7.9 mark. So, we estimate the length as 0.05 cm



more than 7.8 cm. All of these have meaning and are therefore

significant. We express the reading as 7.85 cm, accurate to three significant figures. All of these figures, 7.85, can be used in calculations. In reading B we see that 9.2 cm is definitely known. We can include one estimated digit in our reading, and we estimate the next digit to be zero. Our reading is reported as 9.20 cm. It is accurate to three significant figures.

Rules for Zeros

If a zero represents a measured quantity, it is a significant figure. If it merely locates the decimal point, it is not a significant figure.

Zero Within a Number. In reading the measurement 9.04 cm, the zero represents a measured quantity, just as 9 and 4, and is, therefore, a significant number. A <u>zero between any of the other digits in a numberisasignificant figure</u>.

Zero at the Front of a Number. In reading the measurement 0.46 cm, the zero does not represent a measured quantity, but merely locates the decimal point. It is not a significant figure. Also, in the measurement 0.07 kg, the zeros are used merely to locate the decimal point and are, therefore, not significant. Zerosatthefirst(left)ofanumberarenotsignificantfigures.

Zero at the End of a Number. In reading the measurement 11.30 cm, the zero is an estimate and represents a measured quantity. It is therefore significant. Another way to look at this: The zero is not needed as a placeholder, and yet it was included by the person recording the measurement. It must have been recorded as a part of the measurement, making it significant. Zerostotherightofthedecimalpoint, and attheendofthenumber, are significant figures.

Zeros at the End of a Whole Number. Zeros at the end of a whole number may or may not be significant. If a distance is reported as 1600 feet, one assumes two sig figs. Reporting measurements in scientific notation removes all doubt, since all numbers written in scientific notation are considered significant. 1 600 feet 1 600 1.6×10^3 feet 1.60×10^3 feet significant figures Four

feet 1.600 x 10³ feet significant figures

1 600 feet

Two significant figures Three

Sample Problem #1: Underline the significant figures in the following numbers. (a) 0.0420 cm answer = 0.0420 cm (e) 2 403 ft. answer = 2403 ft. (b) 5.320 in. answer = 5.320 in. (f) 80.5300 m answer = 80.5300 m

(c) (d) answer = $\underline{10}$ lb. answer = $\underline{2.4}$ x = $\underline{10}$ lb. answer = $\underline{2.0}$ g answer = $\underline{2.4}$ x = $\underline{10}$ lb. answer = $\underline{10}$

Rounding Off Numbers

In reporting a numerical answer, one needs to know how to "round off" a number to include the correct number of significant figures. Even in a series of operations leading to the final answer, one must "round off" numbers. The rules are well accepted rules:

- 1. If the figure to be dropped is less than 5, simply eliminate it.
- 2. If the figure to be dropped is greater than 5, eliminate it and raise the preceding figure by
- 1. 3. If the figure is 5, followed by nonzero digits, raise the preceding figure by 1
- 4. If the figure is 5, not followed by nonzero digit(s), and preceded by an odd digit, raise the preceding digit by one
- 5. If the figure is 5, not followed by nonzero digit(s), and the preceding significant digit is even, the preceding digit remains unchanged

Sample Problem #2: Round off the following to three significant figures.

(a) 3.478 m answer = 3.48 m (c) 5.333 g answer = 5.33 g (b) 4.8055 cm answer = 4.81 cm (d) 7.999 in. answer = 8.00 in.

<u>Multiplication</u>

In multiplying two numbers, when you wish to determine the number of significant figures you should have in your answer (the product), you should inspect the numbers multiplied and find which has the least number of significant figures. This is the number of significant figures you should have in your answer (the product). Thus the answer to 0.024×1244 would be rounded off to contain two significant figures since the factor with the lesser number of significant figures (0.024) has only two such figures.

Sample Problem #3: Find the area of a rectangle 2.1 cm by 3.24 cm.

```
Solution: Area = 2.1 \text{ cm x } 3.24 \text{ cm} = 6.804 \text{ cm}^2
```

We note that 2.1 contains two significant figures, while 3.24 contains three significant figures. Our product should contain no more than two significant figures. Therefore, our answer would be recorded as 6.8 cm² Sample Problem #4: Find the volume of a rectangular solid 10.2 cm x 8.24 cm x 1.8 cm Solution: Volume

```
= 10.2 \text{ cm x } 8.24 \text{ cm x } 1.8 \text{ cm} = 151.2864 \text{ cm}^3
```

We observe that the factor having the least number of significant figures is 1.8 cm. It contains two significant figures. Therefore, the answer is rounded off to 150 cm³.

Division

In dividing two numbers, the answer (quotient) should contain the same number of significant figures as are contained in the number (divisor or dividend) with the least number of significant figures. Thus the answer to 528 © 0.14 would be rounded off to contain two significant figures. The answer to 0.340 © 3242 would be rounded off to contain three significant figures.

```
Sample Problem #5: Calculate 20.45 ② 2.4 Solution: 20.45 ② 2.4 = 8.52083
```

We note that the 2.4 has fewer significant figures than the 20.45. It has only two significant figures. Therefore, our answer should have no more than two significant figures and should be reported as 8.5.

Addition and Subtraction

In adding (or subtracting), set down the numbers, being sure to keep like decimal places under each other, and add (or subtract). Next, note which column contains the first estimated figure. This column determines the last decimal place of the answer. After the answer is obtained, it should be rounded off in this column. In other words, round to the least number of decimal places in you data.

```
Sample Problem #6: Add 42.56 g + 39.460 g + 4.1g
Solution:
42.56 g
39.460 g
4.1g
```

Sum = 86.120 g

Since the number 4.1 only extends to the first decimal place, the answer must be rounded to the first decimal place, yielding the answer 86.1 g.

Average Readings

The average of a number of successive readings will have the same number of decimal places that are in their sum.

Sample Problem #7: A graduated cylinder was weighed three times and the recorded weighings were 12.523 g, 12.497 g, 12.515 g. Calculate the average weight.

Solution:

12.523 g 12.497 g 12.515g 37.535 g

In order to find the average, the sum is divided by 3 to give an answer of 12.51167. Since each number extends to three decimal places, the final answer is rounded to three decimal places, yielding a final answer of 12.512 g. Notice that the divisor of 3 does not effect the rounding of the final answer. This is because 3 is an exact number - known to an infinite number of decimal places.

SCIENTIFIC NOTATION, SIG. FIGS., DENSITY

1. Change to scientific notation.

a.
$$5.420 \times 10^3 = d. 0.0067 \times 10^{-4} = b. 0.020 \times 10^3 = e. -870 \times 10^{-4} =$$

c.
$$0.00492 \times 10^{12} = f. -602 \times 10^{21} =$$

2. Determine the number of sig. figs. in the following:

a.
$$0.002030 = s.f.$$
 e. $670 = s.f.$ b. $670.0 = s.f.$ f. $1.35000 = s.f.$

$$\frac{2}{4 \times 10^2}$$
 = s.f. c.

$$0.060 \times 10^3 = s.f$$

d.

$$4640 = \text{s.f.}$$

g. $4.00 \times 10^2 = \text{s.f. h.}$

3. Perform the following calculations. Report your answer in correct number of sig.figs. and units.

a.
$$1.008 \text{ m} + 32.00 \text{ m} + 2.2 \text{ m} =$$

b.
$$17.65 \text{ g} - 9.7 \text{ g} =$$

$$c. 2.03 \text{ cm}^2 \div 1.2 \text{ cm} =$$

d.
$$13.8612 \text{ cm x } 2.02 \text{ cm} =$$

4. Givethe number of significant figures in each of the following:

402 m 34.20 lbs 0.03 sec 0.00420 g 3 200 liters 0.0300 ft.

(5)a Multiply each of the following, observing sig fig rules

$$= 0.050 \text{ m} \times 102 \text{ m} =$$

$$0.005 \text{ in x } 8 888 \text{ in} =$$

$$324\ 000\ \text{cm}\ \text{x}\ 12.00\ \text{cm} =$$

0.424 in x .090 in =

17 m x 324 m =

5b. Divide each of the following, observing sig figs rules

$$23.4 \text{ m} \div 0.50 \text{ sec} = 0.960 \text{ g} \div 1.51 \text{ moles} =$$

12 miles
$$\div$$
 3.20 hours =

$$1\ 200\ \text{m} \div 12.12\ \text{sec} =$$

6. Add each of the following, observing significant figures rules:

0.5 m 1.9999g 5.9 cm

7. Subtract each of the following, observing significant figure rules:

42.306 m 14.33 g 234.1 cm -1.22 m -3.468g -62.04cm

8. Work each of the following problems, observing sig fig rules

- a) Three determinations were made of the percentage of oxygen in mercuric oxide. The results were 7.40%, 7.43%, and 7.35%. What was the average percentage?
- b) A rectangular solid measures 13.4 cm x 11.0 cm x 2.2 cm. Calculate the volume of the solid. c)

If the density of mercury is 13.6 g/ml, what is the mass in grams of 3426 ml of the liquid?

- d) A copper cylinder, 12.0 cm in radius, is 44.0 cm long. If the density of copper is 8.90 g/cm^3 , calculate the mass in grams of the cylinder. (assume pi = 3.14)
- 9. Write at least 7 facts describing/explaining the Chemistry

behind the picture on the right.

10. A 12.00 g unknown substance is placed in a container with 50.0 mL water. The water level rose up to 55.2 mL. Calculate the density of the substance.



11. Draw/illustrate how the following substances will appear inside a graduated cylinder just like in question # 9.

Liquid 1 ($0.69~\mbox{g/mL}$) Solid 1 ($0.8~\mbox{g/mL}$)

Liquid 2 (1.26 g/mL) Solid 2 (2.04 g/mL)

Liquid 3 (water) Liquid 4 (3.05 g/mL)

FORMULA WRITING, NAMING OF COMPOUNDS & BALANCING CHEMICAL EQUATIONS

(Refer to the Periodic Table included in this packet as well as the list of polyatomic ions)

- 12. Name the following ionic compounds: 13. Write ionic formulas for the following compounds:
 - a. LiCl a. sodium acetate -
 - b. $Mg(OH)_2$ b. tin(II) chloride -
 - c. $K_3P c.$ calcium hydroxide -
 - d. $Fe_2O_3 d$. zinc sulfite -
 - e. FeO e. ammonium sulfate -
 - f. ZnCl₂ f. manganese(II) hypochlorite g. AgNO₃ g. copper (I) nitrite -
 - h. NH₄Cl h. silver cyanide -
 - i. CuCl₂ i. sodium chloride –
 - j. SnCl₂ j. lithium fluoride -
 - k. PbO₂ k. potassium sulfide –
 - 1. AlCl₃ 1. aluminum oxide -
 - m. PbSO4 m. nickel (II) chlorite –
 - n. $Mg_3(PO_3)_2 n$. lead (II) nitrate
 - o. Na₂CO₃ o. ammonium sulfate -
 - p. NaHCO₃ p. aluminum perchlorate -
 - g. KCN g. iron (II) dichromate r. KMnO4- r. lead (IV) bromite s. FeC2O4 –
 - s. lead (II) periodate t. Al(ClO)3 t. magnesium thiocyanate u. FeS2O3 u. calcium thiosulfate v. Sn(CrO4)2 v. sodium bicarbonate w. Mg(HSO4)2 w. strontium hydroxide -
 - 14. Name the following covalent 15. Write the molecular formula for compounds: the following compounds:
 - a. CO a. xenon hexafluoride b. CO2 b. tetranitrogen tetraoxide c. H2O c. boron trifluoride d. CCl4 d. carbon tetrabromide e. N2O3 e. dicarbon tetrafluoride f. SiO2 f. nitrogen tribromide g. N2O g. dinitrogen tetrasulfide h. CBr4 h. oxygen difluoride i. SO2 i. dinitrogen pentoxide j. S2Cl2 j. tetraphosphorus decoxide k. P2O7 k. sulfur hexafluoride -
 - 16. Translate the following word equations to a balanced chemical

equations. a. Iron (II) oxide + aluminum iron + aluminum oxide

b. Hydrochloric acid + sodium hydroxide water + sodium chloride c.

Calcium phosphate + sulfuric acid calcium sulfate + phosphoric acid

d. calcium carbonate calcium + carbon + oxygen gas

e. sodium chloride + silver nitrate | sodium nitrate + silver chloride

f. potassium hydroxide _+ sulfuric acid potassium sulfate + water 17 .Identify the type of equation for each of the equations you balanced in #16

MOLES ↔ GRAMS, MOLARITY, AND STOICHIOMETRY

- a. Use the Periodic Table included in this packet for the atomic masses. Do not round the atomic masses.
- b. Show cancellation of units and report the final answer with the correct unit and correct number of sig figs.
- c. Molarity is a measurement of concentration. M = mol/L. If the concentration of a substance is 1.5 M, it means that 1.5 mol = 1 L.
- 18. Convert the following to moles:

a.
$$36.85 \text{ g C} =$$

b. 170 g
$$O_2$$
=

c.
$$24.0 \text{ g Cu} =$$

d.
$$165.02 \text{ g H}_2\text{O} =$$

e.
$$320.0 \text{ g CaCO}_3 =$$

f.
$$50.020 \text{ g Ca}_3(PO_4)_2 =$$

19. Convert the following to grams:

a.
$$1.20 \text{ mol H}_2 =$$

- b. 0.052 mol Ca =
- c. $10.0 \text{ mol CO}_2 =$
- d. $0.00650 \text{ mol AgNO}_3 =$
- e. 1.025 mole $Al_2(SO_4)_3 =$
- 20. Find the concentration for each of the following
 - a. 25.0 g HCl in 2.1 L solution
 - b. 42.2 g KOH in 250 mL solution
 - c. 0.065 kg Ba(OH)2 in 350 mL solution
- 21. Find the number of moles of solute present in the following

solutions: a. 1.20 L of 0.25M H2SO4 solution

- b. 0.520 L of 1.2M CuSO4 solution
- c. 650.0 mL of 0.21M KNO3 solution
- 22. Solve the following stoichiometric problems completely.
- a. Air bags in cars operate according to the reaction:

$$-\frac{Na_{3}^{N}}{3(s)}$$
 $-\frac{Na_{(s)}^{+}}{(s)}$ $-\frac{N}{2}(g)$

How many grams of nitrogen gas are produced during the decomposition of

- 3.25 g Na3N?
- b. How many grams of lithium are needed to produce 45.0 g of lithium nitride, according to the following process?

$$-\frac{\text{Li}}{(s)}^{+} - \frac{N}{2(g)} - \frac{\text{Li}}{3}^{N}(s)$$

c. A 24.5g sample of sodium chloride reacts with 41.3 g of fluorine gas according to the following chemical equation:

$$_NaCl(s) + _F_2(g)$$
 $_NaF(s) + _Cl_2(g)$

i. Which is the limiting reactant? Justify your answer with calculations.

ii. How many grams of chlorine gas are produced?

- iii. How much excess is left over?
- d. An 84.1 gram sample of phosphorus reacts with 85.0 g of oxygen according to the following equation:

$$-\frac{P}{(s)}^{+}-\frac{O}{2(g)}^{-}-\frac{P}{2}\frac{O}{5(s)}$$

i. Find the limiting reactant. Justify your answer with calculations.

ii. How many grams of P2O5 are produced in theory ? (based on calculation)
iii. A student performed the reaction in the lab and found out that only 123 g of P2O5 were produced. What then is the percent yield for P2O5 ?
GAS LAWS
23. Write down what the following gas laws state and their respective equations.
a. Boyle's Law : Equation :
b. Charles' Law : Equation :
c. Gay-Lussac's Law : Equation :
d. Combined Gas Law : Equation :
e. Avogadro's Law : Equation :
f. Ideal Gas Law : Equation :
24. A 100.0 L sample of gas was compressed to 10.0 mL where its pressure is 350.0 torr. What was the original pressure (in torr) of the 100.0 L sample?

25. Butane gas is stored in a tank at a pressure of 10.0 atm at 22.0°C. The tank can hold a pressure of 50.0 atm before bursting. During a fire the gas is heated to 500.0°C. What is the gas pressure, and will

the tank contain the gas without bursting?

- 26. Calculate the volume in liters of 38.8 g of CO2 at 725 torr and 25.0°C
- 27. A gas occupies 450.0 mL at 655 mm Hg pressure and 30.0°C. What will its volume be at STP?

28. On hot days, you may have noticed that potato chip bags seem to "inflate", even though they have not been opened. If I have a 250.0 mL bag at a temperature of 19.0° C, and I leave it in my car which has a temperature of 60.0° C, what will the new volume of the bag be?

General problems

- 29. How many grams of Na CO $_2$ (molar mass = 106.0 g/mol) are required for complete reaction with 25.0 mL of 0.155 M HNO ? (hint: M = mol/L; so 0.155 M is equal to 0.155 mol HNO $_3$ = 1 L HNO $_3$ Na CO $_3$ + 2HNO $_3$ $_3$ 2NaNO $_3$ + CO $_3$ + HNO $_3$ $_3$ 2NaNO $_3$ + CO $_3$ + HNO $_3$ 2NaNO $_3$ 2N
- 30. At STP, it was found that 1.12 L of a gas had a mass of 2.78 g. What is it's molar mass? 31. A sample

of gas occupies 30.0 L at 0.800 atm and 298 K. How many moles of gas are in the sample?

32. What volume of 0.150 M NaOH is needed to react completely with 3.45 g iodine according to the equation: ${}^3I_2 + 6 \text{ NaOH} \rightarrow 5 \text{ NaI} + \text{NaIO}_3 + {}^3H_2\text{O}$

33. 1.056 g of metal carbonate, containing an unknown metal, M, were heated to give the metal oxide and 0.376 g CO 2.

$$MCO_3(s) + heat \rightarrow MO(s) + CO_2(g)$$

What is the identity of the metal M? (show all work)

- a) Mg c) Zn
- b) Cu d) Ba
- 34. The reaction of 25.0 g benzene, C $_{6}^{H}$, with excess HNO $_{3}^{T}$ resulted in 21.4 g C $_{6}^{H}$ NO $_{2}^{N}$. What is the percentage yield?

$$\overset{\text{C H}}{_{6}} \overset{+}{_{6}} \overset{\text{H NO}}{_{3}} \overset{-}{_{6}} \overset{\text{C H NO}}{_{5}} \overset{+}{_{2}} \overset{\text{H}}{_{2}} \overset{\text{O}}{_{2}}$$

- 35. A 100.0-g sample of a compound is made up of 35.9 g of aluminum and 64.1 g of sulfur. The empirical of the compounds is:
- 36. An organic compound which has the empirical formula CHO has a molar mass of 232. Determine the molecular formula

37.

How many significant digits are present in the temperature read from the thermometer illustrated to the right?

- b) 356 mg to g
- c) 0.00968 nL to hL
- d) 0.00023480 ks to cs e) 0.000589 nm

to m

- 38. Convert
- a) 23.6 cm to km

a) 1 b) 2 c) 3 d) 4

		Name	_
		Date	
	Saf	ety Quiz	
Fill	in the blank.		
1.	Pour chemicals from large reagent bot	les into	
	before mean	uring.	
2.	Read anda	chemical label before using the chemical.	
3.	When diluting an acid, always add	to	
4.	Work with volatile chemicals under a	×	
5.	Use a or	to draw liquid into a pipette.	
6	Strike matches	your body.	

7.	Move test tubes back and forth at anv	vhile heating.
8.	Hold hot glassware with or	
9.	When inserting lubricated glass tubing into a stopper, protect you	r hands with
	Always protect your eyes with working in the laboratory.	when
11.	Stand on a	if you need to reach.
12.	Rinse chemicals from your eyes in an	
13.	clothing on the way to the safety show	rer.
14.	Extinguish small fires in containers by	them.
15.	Put out clothing fires in a	·

Safety Quiz

1.	Hold bottles with your hand over the label while pouring.
2.	Immediately wipe up any spills.
3.	When lighting a Bunsen burner, first turn on the gas, then strike your match.
4.	Check glassware for stars and cracks.
5.	Shake laboratory thermometers down before use.
6.	Carefilly scoop up mercury from a broken thermometer with a piece of paper.
7.	If you don't have an inserter, lubricate glass tubing before inserting it through stopper.
8.	Balance a centrifuge by placing testo bes containing equal masses opposite each other.
9.	Wear closed leather shoes to protect your feet.

Laboratory Glassware

Name of equipment	Picture	Function (Mark an x to all that apply)						
		Mix	Deliver	Measure	Contain	Deliver	Crush	Η
Beaker								
Thermometer								
Erlenmeyer flask								

Graduated cylinder				
Bunsen burner				
Test tube				
Balance scale				

Name of equipment	Picture	Function (Mark an x to all that apply)							
		Mix	Deliver	Measure	Contain	Deliver	Crush	Н	
Burette (buret)									
Scoopula									
Plastic pipette									
Graduated pipette									
Volumetric flask									
Watchglass									
Mortar and pestle									

Funnel				